

Interrelation of Superconducting and Antiferromagnetic Gaps in high- T_C Compounds: a Test Case for a Microscopic Theory

Werner Hanke^{1,2}, Marc G. Zacher¹, Enrico Arrigoni¹, Shou-Cheng Zhang²

¹*Institute for Theoretical Physics, University of Würzburg, 97074 Würzburg, Germany*

²*Department of Physics, Stanford University, Stanford, California 94305, USA*

Recent angle resolved photoemission (ARPES) data, which found evidence for a d -wave-like modulation of the antiferromagnetic gap, suggest an intimate interrelation between the antiferromagnetic insulator and the superconductor with its d -wave gap. This poses a new challenge to microscopic descriptions, which should account for this correlation between, at first sight, very different states of matter. Here, we propose a microscopic mechanism which provides a definite correlation between these two different gap structures: it is shown that a projected SO(5) theory, which aims at unifying antiferromagnetism and d -wave superconductivity via a common symmetry principle while explicitly taking the Mott-Hubbard gap into account, correctly describes the observed gap characteristics. Specifically, it accounts for both the dispersion and the order of magnitude difference between the antiferromagnetic gap modulation and the superconducting gap.

Especially due to recent advances in experimental techniques, such as angular-resolved photoemission spectroscopy (ARPES) [1–6] and spectroscopy probing magnetic correlations [7,8], a crucial basic ingredient seems to emerge for the phenomenology of the high- T_C compounds: in contrast to many of the earlier theoretical studies, which started from the highly doped system far away from the antiferromagnetic (AF) insulator, the appropriate starting point seems to be the insulating state, corresponding to half-filling in theoretical models or to the undoped situation in experiment. A prominent recent example, which advocates this point of view strongly, is a photoemission observation by the Stanford group [1]. It found evidence suggesting a direct correlation between the d -wave symmetry of the superconducting (SC) gap and an observed d -wave-like modulation of the AF gap. Further experiments employing magnetic spectroscopy, support this view of an intimate interrelation of the AF insulator and the SC state as well [7]. These experimental developments have given the high- T_C research a new focus and challenge: any microscopic theory has to account for this interrelation between, at first glance, radically different states.

The "gap" structure in the AF phase as found by ARPES experiments in insulating $\text{Ca}_2\text{CuO}_2\text{Cl}_2$ [1] is summarized in Fig. 1. These ARPES data display a d -wave-like, i.e. $|\cos k_x - \cos k_y|$ -like dispersion in the one-electron spectral function $A(\vec{k}, \omega)$ with respect to the lowest energy state at $(\frac{\pi}{2}, \frac{\pi}{2})$. The straight line in Fig. 1 shows the d -wave dispersion function along the edge of the magnetic Brillouin Zone (dashed square in the inset of Fig. 1) with a d -wave "energy gap". The inset of Fig. 1 presents the ARPES data and the "gap" features in a two-dimensional plot: on a line drawn from the center of the Brillouin zone to the experimental points, the distance between these points to the intersection of this line with the AF Brillouin zone gives the value of the "gap" at the $\vec{k}$ -point considered. The data closely follow the

$|\cos k_x - \cos k_y|$ -dispersion, depicted in the d -wave full line. This " d -wave" like gap is a modulation of the uniform (s -wave) Mott-Hubbard gap of the order of $U \sim eV$ of the insulating state. Therefore, throughout this paper, we shall use the notation $s + |d|$ to characterize the full gap found in the AF insulator. The crucial point is that these photoemission data suggest that the $|d|$ component of the AF gap in the insulator is also the underlying reason for the celebrated pseudo-gap in the underdoped regime: this "high-energy" pseudo-gap of the order $J \sim 0.1eV$ continuously evolves out of the insulating feature, as documented not only by the same energy scale but again by the same d -wave dispersion [3]. Since, on the other hand, this high-energy feature is closely correlated to the superconducting gap as a function of both doping and momentum [3,4,9], we finally arrive at a crucial constraint on the microscopic theory: such a theory should be able to explain the interrelation between the superconducting gap and the AF gap modulation. Theoretical concepts for the pseudo-gap range from descriptions based on preformed pairs [10], schemes employing resonating valence bond (RVB) singlet formation [11,12] and spin-charge separations [13] to damped spin-density waves [14]. However, the microscopic origin of the pseudo-gap and, in particular, its interrelation with the insulating gap features are still unestablished.

In this report, we shall argue that there exists a theory, which provides rather naturally such an interrelation. This is the SO(5) theory of high-temperature superconductivity which aims at unifying AF and SC via a symmetry principle [15–22]. The symmetry enables the construction of a five-dimensional "superspin" composed out of the three-component AF order parameter and the two-component SC order parameter. An SO(5) "rotation" maps the AF microscopic state and its order parameter onto the superconducting counterpart and vice versa and, thereby, provides a definite prediction of the gap characteristics of both states of matter: we shall demonstrate that a d -wave SC gap and the $|d|$ compo-

nent of the AF gap arise from the same physical origin and can be obtained from each other by an appropriate SO(5) rotation. The obtained AF gap modulation has a characteristic sharp cusp around its minimum at $(k_x, k_y) = (\pm\frac{\pi}{2}, \pm\frac{\pi}{2})$, consistent with the available experimental data [23]. Furthermore, the size of the AF gap modulation and the SC gap are also in good agreement with the experimental values.

Besides finding a modulation of the gap in the AF insulator, the recent ARPES experiment [1] also found a remnant Fermi surface of the AF insulator. This shows that it is appropriate to think of the AF insulator in terms of a condensate of *magnons* on top of a Fermi-liquid like state, just like a superconductor can be viewed as a condensate of *Cooper pairs* on top of a Fermi-liquid state [24]. To be more specific, the variational wave function of an AF insulator with Néel vector pointing in α -direction is given by

$$|\Psi_{AF}\rangle \sim \prod_k (\tilde{u}_k + \tilde{v}_k c_{k+Q}^\dagger \sigma_\alpha c_k) |\Omega\rangle, \quad (1)$$

where $c_k^\dagger$ is a two component spinor creation operator for an electron state with wave vector k , $Q = (\pi, \pi)$ is the AF ordering vector, σ_α are the three Pauli spin matrices, $\tilde{u}_k$ and $\tilde{v}_k$ are the variational parameters and $|\Omega\rangle$ is a Fermi-liquid like state at half-filling. A magnon is defined by the operator $N_\alpha(k) = c_{k+Q}^\dagger \sigma_\alpha c_k$, and is formed by a triplet bound state of a particle and a hole pair with respect to a half-filled state. On the other hand, a SC state is described by the following variational wave function

$$|\Psi_{SC}\rangle \sim \prod_k (u_k + v_k c_k \sigma_y c_{-k}) |\Omega\rangle, \quad (2)$$

where u_k and v_k are the variational parameters for the SC state. Here the Cooper pair is defined by the operator $B(k) = c_k \sigma_y c_{-k}$ and is formed by a singlet pair of holes with respect to a half-filled state. We see that these two seemingly different states of matter are actually formally equivalent, if one replaces the magnon operator $N_\alpha(k)$ by a Cooper pair operator $B(k)$. This “replacement” is exactly provided by the SO(5) rotations operator

$$\pi_\alpha = \sum_k g_k c_{k+Q} \sigma_\alpha \sigma_y c_{-k}, \quad (3)$$

where the form factor $g_k = \text{sgn}(\cos k_x - \cos k_y)$ is uniquely determined by the group closure requirement [19,18]. This operator rotates the magnon and Cooper pair operators into each other according to the following equation:

$$[\pi_\alpha, N_\beta(k)] = \delta_{\alpha\beta} g_k B(k) \quad , \quad [\pi_\alpha^\dagger, B(k)] = g_k N_\alpha(k) \quad (4)$$

The internal wave functions of the elementary constituents of the magnon and Cooper pairs are described the symmetries of $\tilde{u}_k \tilde{v}_k$ and $u_k v_k$ respectively. From the above equation we see immediately that a $d =$

$\cos k_x - \cos k_y$ form of the Cooper pair wave function will translate into a $|d| = |\cos k_x - \cos k_y|$ form of the magnon wave function. This formal equivalence between a magnon condensate and a Cooper pair condensate can be elevated to an SO(5) symmetry principle whose dynamical consequence can be tested both numerically and experimentally. Experimentally, the neutron resonance mode in the SC state can be identified with the pseudo Goldstone modes of the enlarged SO(5) symmetry group, which is a quantum fluctuation from the SC state towards the AF state [15,21]. This identification has recently been used to give an estimate for the condensation energy from the normal to the SC state [21]. Numerically, we have performed exact diagonalization studies of the Hubbard and the $t - J$ models [16,17]. In particular, for the $t - J$ model on a finite sized cluster, the ground state at half-filling is a total spin singlet state. Low energy *bosonic* excitations above this ground state are either magnon or hole pair states. The relative energies of a single magnon and a hole pair states can always be made degenerate by tuning the chemical potential to a special value $\mu = \mu_c$. However, what is surprising is that the multi-magnon and hole pair states also become nearly degenerate at this special value of the chemical potential [17]. This is an indication that the interaction between the magnons and the hole pairs are the nearly the same, originating from a common type of physical interaction. Because of this near degeneracy of the multi magnon and hole pair states, it is argued that the low energy *bosonic* excitations of the $t - J$ model near half-filling can be organized into SO(5) symmetry multiplets [17,20].

The exact SO(5) symmetry [18,19] requires charge excitations at half-filling to have the same gap as the collective spin-wave excitations. This condition is violated in a Mott-Hubbard insulator, which has a large gap ($\sim eV$) to all charge excitations while the spin excitations display no gap. In particular, in an exactly SO(5) symmetric description [18,19], the superconducting gap with its nodes would directly be mapped onto an AF gap, which then would have precisely the same magnitude and would go to zero at the nodes $(\pm\pi/2, \pm\pi/2)$. Taking the Mott-Hubbard gap into account amounts to properly projecting out these “high-energy” processes of order $\sim eV$ (Gutzwiller constraint) in the “low-energy” SO(5) rotation between AF and dSC states. Recently, such a projection procedure has been carefully defined for the bosonic excitations and it was shown that the Gutzwiller constraint can be implemented analytically and exactly within the SO(5) theory [25]. The central hypothesis of the SO(5) theory states that the high T_c superconductors can be described by a fully SO(5) symmetric Hamiltonian supplemented by the Gutzwiller constraint. The Gutzwiller constraint associated with the large on-site interaction reduces the full SO(5) symmetry to a projected SO(5) symmetry with well defined characteristics. The corresponding projected SO(5) model describes the

low-energy bosonic degrees of freedom near the AF/SC transition [25].

In this paper, we shall explore the consequence of the projected SO(5) symmetry in the *fermionic* excitations of the AF and SC states. In particular, we shall argue for the following two main points: (i) The ARPES experiments show that the fermionic quasi-particles have a $s + |d|$ gap structure in the AF state. The projected SO(5) symmetry introduces the s component of the AF gap associated with the large on-site Coulomb energy, which is absent in the case of pure SO(5) symmetry, and quantitatively relates the remaining $|d|$ component of the AF gap with the d -wave SC gap. While the d -wave SC gap is of the order of $J/10$, the obtained AF gap modulation is of the order of J . This is an unexpected consequence of the projection. The correspondence between the $|d|$ component of the AF gap structures and the d -wave SC gap structure is the signature of the projected SO(5) symmetry in the fermionic excitation spectra. (ii) The $t - J$ model with nearest neighbor hopping only gives an s gap structure in the AF state without the $|d|$ gap modulation. Therefore, while the *bosonic* excitation spectra of the $t - J$ model can be organized by the projected SO(5) symmetry, its *fermionic* quasi-particle spectra are *not* compatible with the projected SO(5) symmetry and *not* compatible with the ARPES experiments. On the other hand, microscopically SO(5) symmetric models with proper Gutzwiller projection can describe both the bosonic and the fermionic excitation spectra of the high T_c systems.

Let us first see how a microscopically SO(5) symmetric Hamiltonian with Gutzwiller projection can give rise to the $s + |d|$ gap structure as observed in the ARPES experiments. This Hamiltonian contains the manifestly SO(5)-symmetric terms,

$$H_{\text{kin}} + H_{\text{int}} = \sum_{p,\sigma} \varepsilon_p c_{p,\sigma}^\dagger c_{p,\sigma} + V \sum_{\vec{r}_1, \vec{r}_2} \left\{ \vec{m}(\vec{r}_1) \cdot \vec{m}(\vec{r}_2) + \frac{1}{2} (\Delta(\vec{r}_1) \Delta(\vec{r}_2)^\dagger + \Delta(\vec{r}_1)^\dagger \Delta(\vec{r}_2)) \right\}. \quad (5)$$

Here, H_{kin} stands for the kinetic energy part with band dispersion $\varepsilon_p = -2t(\cos k_x + \cos k_y)$, valid for a nearest-neighbor tight-binding model with hopping amplitude t ($c_{p,\sigma}^\dagger$ creates a hole with momentum p and spin σ). Both H_{kin} and H_{int} comprise the explicitly SO(5)-symmetric part of the Hamiltonian [18,19]. H_{int} contains a spin-spin interaction and a pair-hopping term and, thus, serves as an example of the possible AF $\longleftrightarrow$ dSC rotation, the implications of which on the gap structure we shall investigate below. $\vec{m}(\vec{r}_1)$ and $\Delta(\vec{r}_1)$ are Néel and d -wave order parameters (operators) at site $\vec{r}_1$. The d -wave order parameter is built up of the usual superposition of nearest-neighbor (n.n.) d -wave pairs [26]. On the other hand, the Néel order parameter has an extended internal structure [27]. This extended internal structure in $\vec{m}(\vec{r}_1)$

is required by the SO(5) symmetry, which may, at least in principle, be tested in experiments. In particular, it may be related effectively to spatially extended hoppings t' and t'' [27]. The essential physics of H_{int} is that it is embedded, via the spin interactions and pair hopping processes, the dynamical equivalence of spin (triplet or magnon) excitations and pair excitations. We will argue below in more detail that this specific choice is physically motivated in that it provides a possible AF $\longleftrightarrow$ d-SC rotation, when the SC-part in H_{int} corresponds to the usual BCS reduced Hamiltonian for n.n. d -wave pairing. In particular, it will be shown that the gap modulation is introduced in a rather natural way already at the simplest mean-field level, i.e. in terms of BCS and spin-density wave (SDW) gap equations (Hartree-Fock).

In addition to the fully SO(5) symmetric Hamiltonian, we need to implement the Gutzwiller projection, which will reduce the full SO(5) symmetry to a projected SO(5) symmetry. This can be implemented by the introduction of a Hubbard U interaction and by taking the limit of large U . Therefore, we arrive at the following Hamiltonian

$$H = (H_{\text{kin}} + H_{\text{int}}) + H_U + H_\mu, \quad (6)$$

where

$$H_U = U \sum_i n_{i\uparrow} n_{i\downarrow}. \quad (7)$$

($n_{i\uparrow} = c_{i\uparrow}^\dagger c_{i\downarrow}$) is the standard Hubbard interaction and $H_\mu = -\mu Q$ denotes the chemical potential term. At this stage, a cautious reader may wonder about the problem of double counting, since the perturbation theory of $H_{\text{kin}} + H_U$ may produce terms already included in H_{int} . But these perturbation effects are of the order of t^2/U . Therefore, if we can properly take the $U \rightarrow \infty$ limit, we do not have to worry about the double counting problem. There are various ways to (approximately) study the Hamiltonian in Eq. (6). Its physical content becomes transparent already on the simplest, i.e. Hartree-Fock mean-field level. Earlier work by Schrieffer et al. on the Hubbard model [28] shows that such a simple mean field calculation can capture the basic physics in the strong-coupling limit. Consider first the SDW-type of solution for the Néel state. Here the gap function $\Delta(\vec{p})$ is connected to the SDW mean-field (polarized in z -direction) by the standard relation:

$$\langle c_{\vec{p}+\vec{Q},i}^\dagger \sigma_{ij}^3 c_{\vec{p},j} \rangle = \frac{\Delta(\vec{p})}{2E(\vec{p})}, \quad (8)$$

where, as usual, $E(\vec{p}) = (\varepsilon^2(\vec{p}) + \Delta(\vec{p})^2)^{1/2}$. When introduced in equation (6) for the Hamiltonian, this mean-field order parameter results in the self-consistency condition determining the gap $\Delta(\vec{p})$,

$$\sum_{\vec{p}'} V(\vec{p}, \vec{p}') \frac{\Delta(\vec{p}')}{2E(\vec{p}')} = \Delta(\vec{p}). \quad (9)$$

Here we shall take the factorized form of the SO(5) interaction, i.e. $V w_{\vec{p}} w_{\vec{p}'}$ [26], which also follows from equation (5) and the Fourier-transformed expression for the order parameter $\vec{m}(\vec{r})$ in Ref. [27]. Including the Hubbard U -term, $V(\vec{p}, \vec{p}')$ is given by

$$V(\vec{p}, \vec{p}') = U + V w_{\vec{p}} w_{\vec{p}'}, \quad (10)$$

It is now straightforward to show that the gap equation (9) for the Néel state as well as the corresponding BCS-type gap equation for the SC state yield the required gap features:

First, note that the factorized form of the interaction $V(\vec{p}, \vec{p}')$ introduces a similarly separable form of the AF gap,

$$\Delta^{\text{AF}}(\vec{p}) = \Delta_U + \Delta_{\text{mod}} w_{\vec{p}}, \quad (11)$$

where, as before [26], $w_{\vec{p}} = |\cos p_x - \cos p_y|$. For large values of the Hubbard interaction Δ_U is of the order of U . From experiment, Δ_{mod} should be of order J while Δ_{SC} should be of the order of $J/10$ (this will be shown below to be the case using a Slave-Boson mean-field evaluation). Equation (11) then establishes the gap-modulation $\sim J|\cos p_x - \cos p_y|$ on top of a uniform gap in the AF (Fig. 3).

Second, in formal analogy, in the d -wave SC state, the same gap equation (9) holds, with three modifications: The first one is, of course, that $\Delta^{\text{SC}}(\vec{p})$ stands now for the d -wave SC order parameter with n.n. pairing [26]. The second is due to the well-known fact, that the U -term and, therefore, the constant gap term of the AF drops out. This is because of the condition $\sum_{\vec{p}} U \frac{\Delta^{\text{SC}}(\vec{p})}{2E(\vec{p})} = 0$, which in real space is nothing but the two pairing electrons avoiding the on-site Coulomb repulsion [29].

Third, in the SC state, the relevant interaction in the gap equation is $V(\vec{p}, \vec{p}') = V (w_{\vec{p}} \text{sgn}(\cos p_x - \cos p_y)) \cdot (w_{\vec{p}'} \text{sgn}(\cos p'_x - \cos p'_y))$, finally resulting in the gap function (Fig. 4),

$$\Delta^{\text{SC}}(\vec{p}) = \Delta^{\text{SC}} \cdot (\cos p_x - \cos p_y). \quad (12)$$

Thus, both the AF gap in equation (11) and the SC gap have the required form. What experiments tell us is that while Δ^{AF} is of the order J , Δ^{SC} is an order of magnitude smaller. This requires a more involved mean-field evaluation: We have chosen a technique that is capable of dealing with the problem in a way which is non-perturbative in the Hubbard interaction U . It is the Slave-Boson formalism introduced by Kotliar and Ruckenstein [30], which we treat by the usual saddle-point approximation.

Confidence in this approximation derives from various observations: (i) the Slave-Boson mean-field approximation gives the identical result as the so-called Gutzwiller approximation. The latter is equivalent to the exact treatment of the Gutzwiller variational approach in infinite dimensions [30,31]. We shall demonstrate that this result is relevant for the "projected" SO(5) theory, i.e. for projecting out the higher-energy states of the upper Hubbard band, as discussed above. (ii) secondly, this approximation yields rather satisfying agreement with, in principle exact, QMC simulations over a wide range of Coulomb correlations U and values for the doping [32]. (iii) the two main results of our study, i.e. that the AF gap modulation itself is correlated to the SC gap by symmetry and that the Hubbard term H_U induces the order of magnitude difference in the effects should be independent of specific approximations. This will be explicitly verified by comparing with the bare Hartree-Fock (HF) study.

The results of the Slave-Boson evaluation of the Hamiltonian are presented in terms of the gap features in Figs. 3, 4 and 5.

Figs. 3 and 4 display the SC gap and the AF gap modulation again in the same 2D version as in the experimental plot in Fig. 1. We note that as in HF, for symmetry reasons, the d -wave gap in the SC leads to a $|\cos k_x - \cos k_y|$ -modulation of the AF gap. Thus, the AF gap is given by $\Delta^{\text{AF}}(\vec{k}) = \Delta_U + \Delta_{\text{mod}} |\cos k_x - \cos k_y|$ displaying a cusp at the $(k_x, k_y) = (\pm \frac{\pi}{2}, \pm \frac{\pi}{2})$ wave vectors. The theory has only one free parameter, namely the SO(5)-coupling strength V in Eq. (5), which was chosen in such a way that it gives a d -wave gap of the correct order of magnitude in the SC phase, i.e. $\Delta_{\text{SC}} = 0.02t \approx J/10$ [33]. The crucial observation in Figs. 3, 4 and 5 is then that, while the SO(5) interaction is responsible for the d -wave structure of both gaps, it is a different mechanism, namely the Hubbard gap, which is responsible for the experimentally observed order of magnitude differences in $\Delta_{\text{SC}} \approx J/10$ and the AF gap modulation $\Delta_{\text{mod}} \approx J$.

This is clearly demonstrated in Fig. 5, which plots the amplitude Δ_{mod} of the d -wave-like modulation as a function of the Hubbard-gap Δ_U (Δ_U scales with U for large U). We note that increasing U (and, thus, Δ_U) and, therefore, projecting out the doubly occupied states, strongly enhances the d -wave-like modulation in the AF gap. In particular, taking commonly accepted values $U = 8t$ and, therefore, $\Delta_U \approx 3t$ [34,35] yields an AF gap modulation of order $\Delta_{\text{mod}} \approx (0.25 - 0.3)t$. Thus, we find a radically different energy scale for Δ_{mod} of order J and Δ_{SC} of order $J/10$, in agreement with the ARPES data.

Let us now comment on the possible alternative explanations of the ARPES data. Both Hubbard and t - J Hamiltonians have been extensively studied in two dimen-

sions by employing exact diagonalization and Quantum-Monte-Carlo (QMC) numerical techniques (see, for example, Refs. [34–37]). There is general consensus that both models have a serious deficiency, when compared with the ARPES data in Fig. 1: they do not display the d -wave-type dispersion, and the energies at $(\pi, 0)$ and $(\frac{\pi}{2}, \frac{\pi}{2})$ are essentially degenerate [34–36]; as we have mentioned above, this is serious, since ARPES shows that the evolution of the $(\pi, 0)$ feature is crucial to understand the d -wave-like pseudo-gap. These numerical “experiments” have been further carried out with the inclusion of more-distant (next-nearest t' , t'' etc.) neighbor hoppings [5,36,37].

However, this procedure requires delicate adjustments of the hopping integrals to shift the structure near $(\pi, 0)$ to the observed binding energy. In addition, at first sight, it seems to be completely uncorrelated with two experimental findings which emphasize the universal role of the magnetic energy scale J : one is the fact that the insulating bandwidth itself scales with J [1]. The other is that, at present, it is not clear whether the specific t' , t'' -choice is simultaneously in accordance with the universal picture which recently emerged from the incommensurate spin fluctuations of two major classes of high- T_c materials, $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ and $\text{YBa}_2\text{Cu}_3\text{O}_{7-x}$ [7,8]. We have already indicated (see Ref. [26]) that t' , t'' -hoppings may arise effectively from the $\text{SO}(5)$ -part of the Hamiltonian in Eq. (5), namely from the spin interaction term $V \sum_{\vec{r}_1, \vec{r}_2} \vec{m}(\vec{r}_1) \cdot \vec{m}(\vec{r}_2)$. However, in addition to t' , t'' this interaction effectively produces longer-range hoppings (although with magnitude decreasing with distance) which are essential in order to produce the cusp-like feature of the AF-gap. Indeed, a finite-range hopping (i.e. limited to t' , t'') is not sufficient to produce a cusp-like feature [38].

Summarizing, our results show that the recent experimental discovery of the $s + |d|$ gap structure in the AF state can be naturally explained by the concept of projected $\text{SO}(5)$ symmetry, which relates the $|d|$ gap modulation of the AF state with the d -wave gap of the SC state. If taken as an exact $\text{SO}(5)$ theory, without the physically relevant symmetry-breaking term H_U , this would have resulted in an exact mapping, where the AF gap has nodes. It has been shown here that, while H_U breaks the symmetry and restores the Mott-Hubbard gap, it does not break the experimentally observed correlation between the d -wave gap features. In fact, as demonstrated here, H_U is pivotal in explaining the order of magnitude differences between the superconducting gap Δ_{SC} and the d -wave-like modulation of the AF gap Δ_{mod} . Just like the neutron resonance mode can be interpreted as the reflection of AF correlation in the SC state, the ARPES experiment can be interpreted as the reflection of the SC correlation in the AF state. Our results also show that the Hubbard and $t - J$ models still “miss a piece”, while a microscopically $\text{SO}(5)$ symmet-

ric Hamiltonian with proper Gutzwiller projection can correctly describe both the bosonic and the fermionic excitation spectra of the high T_c superconductors.

We thank R.B. Laughlin, D.J. Scalapino, R. Eder, and O.K. Andersen for helpful discussions. W.H. and E.A. would also like to acknowledge the support and hospitality of the Stanford Physics Department, where most of this work was carried out. This work was supported from the DFG (AR 324/1-1 and HA 1537/17-1), DFN (contract No. TK598-VA/D3), BMBF (05SB8WWA1) and NSF (grant number DMR-9814289).

-
- [1] F. Ronning et al., *Science* **282**, 2067 (1998).
 - [2] Z.X. Shen et al., *Phys. Rev. Lett.* **70**, 1553 (1993); A.G. Loeser et al., *Science* **273**, 325 (1996); B.O. Wells et al., *Phys. Rev. Lett.* **74**, 964 (1995).
 - [3] D.S. Marshall et al., *Phys. Rev. Lett.* **76**, 4841 (1996).
 - [4] H. Ding et al., *Nature* **382**, 51 (1996).
 - [5] C. Kim et al., *Phys. Rev. Lett.* **80**, 4245 (1998).
 - [6] A. Ino et al., *Phys. Rev. Lett.* **81**, 2124 (1998).
 - [7] G. Aeppli et al., *Science* **278**, 1432 (1997); M.A. Kastner et al., *Rev. Mod. Phys.* **70**, 897 (1998).
 - [8] H.A. Mook et al., *Nature* **395**, 580 (1998).
 - [9] P.J. White et al., *Phys. Rev. B* **54**, R15669 (1996).
 - [10] S.A. Kivelson and V.J. Emery, *Nature* **374**, 434 (1995); C.A.R. de sa Melo, M. Randeria, J. Engelbrecht, *Phys. Rev. Lett.* **71**, 3202 (1993); N. Trivedi and M. Randeria, *ibid.* **75**, 312 (1995); S. Doniach and M. Inui, *Phys. Rev. B* **41**, 6668 (1990); A.J. Millis, L.B. Ioffe, H. Monien, J. Phys. Chem. Solids **56**, 1641 (1995); B. Janko, J. Maly, K. Levin, *Phys. Rev. B* **56**, R11407 (1997); Y.L. Uemura et al., *Phys. Rev. Lett.* **62**, 2317 (1989).
 - [11] T. Tanamoto, K. Kohno, H. Fukuyama, *J. Phys. Soc. Jpn.* **61**, 1886 (1992); Y. Suzumura, Y. Hasegawa, H. Fukuyama, *ibid.* **57**, 2768 (1988); G. Kotliar and J. Liu, *Phys. Rev. B* **38**, 5142 (1988).
 - [12] X.-G. Wen and P.A. Lee, *Phys. Rev. Lett.* **78**, 4111 (1997).
 - [13] R.B. Laughlin, *Phys. Rev. Lett.* **79**, 1726 (1997).
 - [14] J. Schmalian, D. Pines, B. Stojkovic, *Phys. Rev. Lett.* **80**, 3839 (1998); J. Schmalian, D. Pines, B. Stojkovic, <http://xxx.lanl.gov/abs/cond-mat/9804129>; A.V. Chubukov and D.K. Morr, <http://xxx.lanl.gov/abs/cond-mat/9806200>.
 - [15] S.C. Zhang, *Science* **275**, 1089 (1997).
 - [16] S. Meixner, W. Hanke, E. Demler and S.C. Zhang, *Phys. Rev. Lett.* **79**, 4902 (1997).
 - [17] R. Eder, W. Hanke and S.C. Zhang, *Phys. Rev. B* **57**, 13781 (1998).
 - [18] S. Rabello, H. Kohno, E. Demler and S.C. Zhang, *Phys. Rev. Lett.* **80**, 3586 (1998).
 - [19] C. Henley, *Phys. Rev. Lett.* **80**, 3590 (1998).
 - [20] R. Eder, A. Dorneich, M.G. Zacher, W. Hanke and S.C. Zhang, *Phys. Rev. B* **59**, 561 (1999).

- [21] E. Demler and S.C. Zhang, *Nature* **396**, 733 (1998).
- [22] E. Arrigoni and W. Hanke, *Phys. Rev. Lett.* **82**, 2115 (1999).
- [23] Notice that the ARPES data measure only the dispersion of the first ionization states, i.e. the "topmost" valence band dispersion. In case of electron-hole symmetry, as in our theory, this dispersion is then translated into an equivalent $|\cos k_x - \cos k_y|$ -modulation of the AF gap.
- [24] While this "normal state" may have anomalous properties it has the same symmetry as the Fermi-liquid state (see Ref. [39]). We identify the normal state as a Fermi-liquid in the sense that it is adiabatically continuable to the over-doped metallic state. Alternatively to this momentum-space description, one may start from a more localized real-space version: above $T_{\text{Néel}}$ up to temperatures of order $J \sim 1000K$, a variety of experiments suggest that the charge carriers have a tendency to bind into spin-singlet states. However, the spin singlets are spatially disordered. They form a spin-liquid state $|\tilde{\Omega}\rangle$, which we may take to be of short-range RVB (resonating-valence-bond) form, where the singlets are formed by nearest-neighbor spins [20,39]. Consider then two "fixed" neighboring sites in the AF, say in the spin configuration $(\uparrow\downarrow)$. This "fixed" configuration can be generated by superposition of the singlet $(\uparrow\downarrow) - (\downarrow\uparrow)$ with the $S_z = 0$ triplet $(\uparrow\downarrow) + (\downarrow\uparrow)$. To obtain the macroscopic 2D antiferromagnet, we have to admix into the spin-liquid at high temperatures a macroscopic number of triplets $t_z(\vec{Q})$. The AF ordered state can then be generated forming the coherent state $|\psi\rangle \sim e^{t_z(\vec{Q})}|\tilde{\Omega}\rangle$, i.e. via a "condensation" of triplets into the lowest-energy state [20,39]. However, in the SO(5) theory, the z-component triplet with momentum (π, π) and the hole pair with momentum $(0, 0)$ are dynamically equivalent, in the same sense as e.g. proton and neutron are considered dynamically equivalent in the isospin theory of nuclear physics. The "RVB type" of spin liquid can thus serve as an alternative common reference state $|\tilde{\Omega}\rangle$ to consider the condensation of magnons and charged Bosons (hole pairs) on equal footing, thus revealing the analogy between AF and d-SC states. The detailed relation between the Fermi-liquid momentum-space description and the real-space spin liquid picture is not clear yet and presently under intensive study.
- [25] S.C. Zhang, J.P. Hu, E. Arrigoni, W. Hanke, and A. Auerbach, <http://xxx.lanl.gov/abs/cond-mat/9904142>.
- [26] The SC part of H_{int} is, in momentum space, $2V\{\Delta\Delta^\dagger + \Delta^\dagger\Delta\}$, where Δ is the usual n.n. d -wave order parameter, i.e. $\Delta = \sum_{p,\sigma} \frac{1}{2}(\cos p_x - \cos p_y)c_{p,\uparrow}^\dagger c_{-\vec{p},\downarrow}$. This "separable" form in Δ and $\Delta^\dagger$ fixes the general SO(5)-invariant interaction with the coupling $V(\vec{p}, \vec{p}'; \vec{q})$ to be separable in momentum space and given by $V(\vec{p}, \vec{p}'; \vec{q}) = V\delta_{\vec{q},\vec{Q}}w_{\vec{p}}w_{\vec{p}'}$, where $w_{\vec{p}} = |\cos p_x - \cos p_y|$ (see also Ref. [18]).
- [27] The extended internal structure of $\tilde{m}(\vec{r}_1)$ [i.e. $\tilde{m}(\vec{r}_1) = \sum_{\vec{r}_2,i,j} w(\vec{r}_1 - \vec{r}_2)e^{i\vec{Q}\cdot\vec{r}_2}c_{\vec{r}_1,i}^\dagger \vec{\sigma}_{ij}c_{\vec{r}_2,j}$, where $\vec{\sigma}$ are the usual 2×2 Pauli matrices] introduces on a mean field level (replacing $\tilde{m}(\vec{r}_2)$ in equation (5) by its expectation value $\langle \tilde{m}(\vec{r}_2) \rangle$) longer-ranged hopping terms t' (next n.n.) and t'' (third n.n.). Similarly to the commonly used spatially extended hoppings [5,36,37], their signs alternate. This stems from the fact that $w(\vec{r})$ is the Fourier-transform of $w(\vec{p}) = |\cos p_x - \cos p_y|$ (see Ref. [26]) and takes the values ($\vec{r} = (m, n)$ on the 2D lattice): $w(m, n) = 0$, if $m+n = \text{odd}$, $w(0, 0) = \frac{8}{\pi^2}$, $w(1, 1) = -\frac{8}{3\pi^2}$ and $w(2, 0) = \frac{8}{9\pi^2}$, etc. This results in $t' \sim -4Vw(0, 0)w(1, 1) \approx -O(J)$ and $t'' \sim +4Vw(0, 0)w(2, 0)$ with $t''/t' = -1/3$, where we used $V = 0.05t$ (see Ref. [33]).
- [28] J.R. Schrieffer, X.G. Wen, and S.C. Zhang, *Phys. Rev. B* **39**, 11663 (1989).
- [29] D. J. Scalapino, *Phys. Rep.* **250**, 329 (1995).
- [30] G. Kotliar and A.E. Ruckenstein, *Phys. Rev. Lett.* **57**, 1362 (1986).
- [31] E. Arrigoni and G. Strinati, *Phys. Rev. Lett.* **71**, 3178 (1993); *Phys. Rev. B* **52**, 2428 (1995); *ibid.* **52**, 13707 (1995).
- [32] See, for example, L. Lilly, A. Muramatsu and W. Hanke, *Phys. Rev. Lett.* **65**, 1379 (1990).
- [33] For the SC gap (around optimal doping) experiments give a d -wave gap magnitude of order $J/10 \sim 10 - 20\text{meV}$ []. A reasonable value for J is $J = 0.2t$. For the choice $V = 0.05t$, we obtain - for the commonly accepted value of the Hubbard correlation $U = 8t$ - at 10 % doping $\Delta_{SC} = 0.02t \approx J/10$.
- [34] For references see E. Dagotto, *Rev. Mod. Phys.* **66**, 76 (1994).
- [35] R. Preuss et al., *Phys. Rev. Lett.* **75**, 1344 (1995); *ibid.* **79**, 1122 (1997).
- [36] R. Eder et al., *Phys. Rev. B* **55**, R3414 (1997).
- [37] D. Duffy et al., *Phys. Rev. B* **56**, 5597 (1997), and references therein.
- [38] There exists preliminary (QMC) numerical evidence that the t' -induced modulation does not follow the linear in $|\cos k_x - \cos k_y|$ -form (along from magnetic BZ), but the SDW-like [see Ref. [1]] ($\sim \cos k_x \cos k_y$)-form: C. Gröber *et al.*
- [39] R. Eder, <http://xxx.lanl.gov/abs/cond-mat/9810216>.




FIG. 1. ARPES results for $\text{Ca}_2\text{CuO}_2\text{Cl}_2$: The lowest energy peak disperses away from the $(\pi/2, \pi/2)$ peak position as one moves away from wavevector $(\pi/2, \pi/2)$. The difference (gap-modulation) between this dispersive peak and the $(\pi/2, \pi/2)$ peak is plotted against $|\cos(k_x a) - \cos(k_y a)|$. The data fit the straight line which indicates a d -wave like dispersion. The inset shows the gap modulation in momentum space along the antiferromagnetic zone boundary. By plotting only the *difference* to $(\pi/2, \pi/2)$, the plot goes to zero for this point by definition.




FIG. 2. The (T, μ) -phase diagram for high- T_c superconductors reveals the intimate relation between the AF and dSC state: The AF state can be viewed as a condensation of magnons (triplet excitations) on top of a Fermi-liquid like state, just as the superconducting state can be represented as the condensation of hole pairs. $\text{SO}(5)$ symmetry relates the magnon onto the hole-pair states. The ARPES experiments probe the internal structure of a Cooper pair and magnon, respectively, thereby revealing the correlation between anti-ferromagnetism and superconductivity.




FIG. 3. Schematic drawing of the gap modulation and the constant gap in the antiferromagnetic phase as obtained from the Slave-Boson Mean-Field calculation. For graphical reasons we indicate the Hubbard gap Δ_U only as a relatively small offset to the modulation. In reality (and in our calculations) $\Delta_U \approx 4t$ is about one order of magnitude larger than the gap modulation. In the analog experimental plot in the inset of Figure 1 only the *difference* to the constant gap is shown and therefore no offset is indicated.




FIG. 4. Schematic drawing of the dSC-gap in the superconducting phase with conventions as in Figure 3 and the inset of Figure 1.




FIG. 5. The key result of the Slave-Boson Mean-Field calculation is the strong enhancement of the gap modulation in the antiferromagnetic phase as U is increased and double occupancy becomes more and more forbidden. The symmetry-breaking U is responsible for a one order of magnitude difference of the gap modulation in the AF phase to the dSC gap in the superconducting phase in the experimentally relevant region of $U \approx 8t$.